arXiv:1208.2021v1 [hep-ph] 9 Aug 2012

Electromagnetic Structure of Pion

Clayton S. Mello, José P. Cruz Filho, Edson O. da Silva, Bruno
El-Bennich, J. P. B. C. de Melo and Victo S. Filho

Laboratório de Física Teórica e Computacional (LFTC),
Universidade Cruzeiro do Sul, 01506-000, São Paulo, Brazil

Abstract. In this work, we analyze the electromagnetic structure of the pion, an elementary particle composed by a quark-antiquark bound state, by considering the calculation of its electromagnetic radius and its electromagnetic form factor in low and intermediate energy range. Such observables are determined by means of a theoretical model that takes into account the constituent quark and antiquark of the pion, in the formalism of the light-front field theory. In particular, it is considered a nonsymmetrical vertex for such a model, in which we have calculated the electromagnetic form factor of the pion in an optimized way, by varying its regulator mass, so that we can obtain the best value for the pion electromagnetic radius when compared with the experimental one. The theoretical calculations are also compared with the most recent experimental data involving the pion electromagnetic form factor and the results show very good agreement.

Keywords: Pion, Electromagnetic Form Factor, Electromagnetic Radius, Light-Front Formalism
PACS: 13.40.Hq, 14.40.Be, 25.80.Ek, 13.40.Gp

INTRODUCTION

In the experimental investigation of pion, many papers have reported a lot of experimental data concerning about its electromagnetic structure, by measuring its electromagnetic form factor $F_\pi(q^2)$, as we can see in Refs. [1, 2, 3, 4, 5, 6, 7]. In the theory of light mesons, a lot of interesting problems has been considered in the literature, as the calculation of the mass and the decay constant of pion in a relativistic potential model of independent quarks [8], the analysis of masses and electroweak properties of light mesons in the relativistic quark model [9] and the study of a relativistic treatment of pion wave functions in the annihilation $\bar{p}p \to \pi - \pi^+$ [10], beside a lot of other works in which are considered calculations involving light mesons, as in Refs. [11, 12, 13, 14, 15].

One of the theoretical ways for describing so many experimental data is to adopt the light-front field theory formalism [16]. A model inspired on QCD was proposed in Ref. [17]. In the present work, we consider in special a description based on a nonsymmetrical vertex model, as early reported in Ref. [18]. In the latter, the light-front formalism is applied to calculate the electromagnetic form factor from currents for quark-antiquark bound states of the pion. Methods like that, based on the light-front formalism, have had an impressive successful in the description of the electromagnetic properties of the hadronic wave functions [17, 18, 19, 20, 21, 22, 23, 24, 25].

In the following, the model considered in the light-front formalism within the nonsymmetrical vertex model for the pion is briefly described. At last, we discuss the numerical results obtained for pion electromagnetic form factor and electromagnetic radius and we discuss the main conclusions.

THE MODEL

The numerical calculation of pion electromagnetic form factor in light-front field theory formalism within the nonsymmetrical vertex model (NSV model) can be performed in the covariant form as:

$$(p+p')^{\mu}F_{\pi}(q^2) \ = <\pi(p')|J^{\mu}|\pi(p)>, \tag{1}$$

in which $q = p' - p$ and the matrix elements of the electromagnetic current $J^{\mu} = e(p^{\mu} + p'^{\mu})F_{\pi}(q^2)$ are given by

$$J^{\mu} = -2\imath e N_c \frac{m^2}{f_{\pi}^2} \int \frac{d^4k}{(2\pi)^4} Tr\left[S(k)\gamma^5 S(k-p')\gamma^{\mu} S(k-p)\gamma^5\right]\Gamma(k,p')\Gamma(k,p), \tag{2}$$

in which $S(p) = (\not{p} - m + \imath\varepsilon)^{-1}$ is the quark propagator and $N_c = 3$ is the number of colors. Such an equation is obtained in the Breit frame, by considering initial momenta $p^{\mu} = (p_0, -q/2, 0, 0)$, final momenta $p'^{\mu} = (p_0, q/2, 0, 0)$ and $q^{\mu} = (0, q, 0, 0)$. As described in previous works [11, 18], the electromagnetic form factor of the pion has only the valence contribution for the plus component of the electromagnetic current. In the case of NSV model, the function $\Gamma(k,p)$ is the regulator vertex function known as nonsymmetrical vertex, that can be written as [11, 18]:

$$\Gamma^{NSY}(k,p) = \frac{N}{(p-k)^2 - m_R^2 + \imath\varepsilon} \ . \tag{3}$$

The pion electromagnetic form factor $F_{\pi}^{(NSY)}(q^2)$ in NSV model, by using the plus component of the electromagnetic current, can be expressed with the light-front wave function, as shown in Refs. [11, 18], according with:

$$F_{\pi}^{(NSY)} = \frac{m^2}{p^+ f_{\pi}^2} N_c \int \frac{d^2k_{\perp} dx}{2(2\pi)^3 x} \mathscr{N}(x,p^+) \Psi_f^{*(NSY)}(x,k_{\perp}) \Psi_i^{(NSY)}(x,k_{\perp}) \theta(x)\theta(1-x), \tag{4}$$

in which $\mathscr{N}(x,p^+) = -4\frac{f_1}{xp^+}(xp^+ - p^+)^2 + 4f_1(xp^+ - 2p^+) + \ xp^+q^2, \ f_1 = k_{\perp}^2 + m^2$ and $x = k^+/p^+$ is the fraction of the carried momentum by the quark. The light-front wave function for the pion with the nonsymmetric vertex can be written as:

$$\Psi^{(NSY)}(x,k_{\perp}) = \frac{N}{(1-x)^2(m_{\pi}^2 - \mathscr{M}_0^2)(m_{\pi}^2 - \mathscr{M}_R^2)}, \tag{5}$$

in which $\mathscr{M}_R^2 = \mathscr{M}^2(m^2, m_R^2) = \frac{k_{\perp}^2 + m^2}{x} + \frac{(p-k)_{\perp}^2 + m_R^2}{(1-x)} - p_{\perp}^2$ and $\mathscr{M}_0^2 = \mathscr{M}^2(m^2, m^2)$ is the free mass operator. The normalization constant N obeys the condition $F_{\pi}(0) = 1$.

NUMERICAL RESULTS

In the numerical calculations, we initially considered the parameters $m_u = m_{\bar{d}} = 0.220$ GeV, $m_R = 0.946$ GeV and $m_{\pi} = 0.140$ GeV. For the input data cited, we obtained for the pion electromagnetic radius the magnitude $\langle r_{\pi^+} \rangle \cong 0.689\, fm$, pointing out an error about 2% in relation to the experimental one, that is, $\langle r_{\pi^+}^{exp} \rangle = (0.672 \pm 0.008)\, fm$ [26]. In order to improve the description of the experimental data for the electromagnetic radius, we explored the variation of the regulator mass m_R of NSV model and we studied its influence on pion electromagnetic form factor for low and intermediate energy range.



FIGURE 1. Plot of the electromagnetic form factor versus $Q^2 = -q^2([GeV/c]^2)$, calculated in NSV model. The different curves correspond to the electromagnetic form factor for some value of regulator mass, that is, m_R=$0.5, 0.8, 0.946, 1.0$ or 1.3 GeV, by keeping constant the light constituent quark masses: m_u=$m_{\bar{d}} = 0.220$ (GeV). The numerical results are compared with the experimental data, as described in Refs. [1, 6] (full triangle up), Ref. [4] (full square), Ref. [5] (full circle), and Ref. [7] (full diamond).

We show in the figure 1 the numerical results obtained from NSV model in the calculation of the pion electromagnetic form factor, in the scale up to 10 $[GeV/c]^2$. By analyzing the curves in the figure 1, we conclude that $m_R = 1.0$ GeV represents the best value for describing with more precision the experimental data, obtained from the references [1, 4, 5, 6, 7]. Values of the regulator mass smaller than $m_R = 0.8$ GeV and above $m_R = 1.3$ GeV do not show good agreement with the measured experimental data for a wide range of energy. In order to confirm the best value of m_R, one can calculate the electromagnetic radius. In fact, we have calculated it for all of the values considered and, for $m_R = 1.0$ GeV, the result obtained, $\langle r_{\pi^+} \rangle \cong 0.673$ fm, represented the best one in relation to the experimental value, with an error less than 0.2%. In order to check our model, we also calculated the pion decay constant for $m_R = 1.0$ GeV, obtaining $f_\pi = 93.1$ MeV, very close to the experimental value $f_\pi^{exp} = 92.2$ MeV.

CONCLUSIONS

We conclude that the best value of the regulator mass for the nonsymmetrical vertex model in the light-front formalism is $m_R = 1.0$ GeV. With that value, it is possible to describe with the best precision the experimental data for the electromagnetic form factor of pion and to obtain the smallest error for the electromagnetic radius in relation to the experimental one. We also studied the dependence of the model on the regulator mass. The numerical results show that the model significantly breaks down for $m_R < 0.8$ GeV and also fails for higher values, above 1.3 GeV, for a wide range of energy.

ACKNOWLEDGMENTS

We thank brazilian agencies CAPES (Coordenadoria de Aperfeiçomento de Pessoal de Nível Superior), CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) and FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo), for financial support.

REFERENCES

1. J. Volmer *et al.*, Phy. Rev. Lett. **86** 1713 (2001).
2. R. Baldini *et al.*, Eur. Phys. J. C 11 **709** (1999).
3. R. Baldini *et al.*, Nucl. Phys. A **666-667** 3 (2000).
4. G. M. Huber *et al.*, Phys. Rev. C **78**, 045203 (2008).
5. V. Tadevosyan *et al.*, Phys Rev. C **75**, 055205 (2007).
6. Brauel *et al.*, Phys. Lett. D **69**, 253 (1977).
7. T. Horn *et al.*, Phys. Rev. Lett. **97**, 192001 (2006).
8. S. N. Jena, M. R. Behera and S. Panda, Pramana J. Phys. **51**, 711 (1998).
9. D. Ebert, R. N. Faustov, and V. O. Galkin, Eur. Phys. J. C **47**, 745 (2006).
10. B. El-Bennich and W. M. Kloet, Phys. Rev. C **70**, 034001 (2004).
11. O. A. T. Dias, Victo S. Filho and J. P. B. C. de Melo, Nuc. Phys. B **199**, 281 (2010).
12. P. Maris and C. D. Roberts, Phys. Rev. C **56**, 3369 (1997).
13. P. Maris, C. D. Roberts and P. C. Tandy, Phys. Lett. B **420**, 267 (1998).
14. A. F. Krutov, V. E. Troitsky and N. A. Tsirova, Phys. Rev. C **80**, 055210 (2009).
15. O. Leitner, J.-F. Mathiot and N. A. Tsirova, Eur. Phys. J. A **47**, 17 (2011).
16. S. J. Brodsky, H. -C. Pauli, and S. S. Pinsky, Phys. Rep. **301**, 299 (1998).
17. T. Frederico and G. Miller, Phys. Rev. D **45**, 4207 (1992).
18. J. P. B. C. de Melo, H. W. L. Naus and T. Frederico, Phys. Rev. C **59**, 2278 (1999).
19. H.-M. Choi and C.-R. Ji, Phys. Rev. D **58**, 071901 (1998).
20. Z. Dziembowski and L. Mankiewicz, Phy. Rev. Lett. **58**, 2175 (1987).
21. F. Cardarelli *et al.*, Phys. Rev. D **53**, 6682 (1996).
22. D. Melikhov and S. Simula, Phys. Rev. D **65**, 094043 (2002).
23. W. Jaus, Phys. Rev. D**60**, 054026 (1999).
24. H. Y. Cheng, C. K. Chua and C. W. Hwang, Phys. Rev. D **69**, 074025 (2004).
25. T. Huang and X. G. Wu, Phys. Rev. D **70**, 093013 (2004).
26. K. Nakamura *et al.* [Particle Data Group Collaboration], J. Phys. G, Nucl. Part. Phys. **37**, 075021 (2010).